

Datasheet: VPA00653KT

Description:	RPL22 ANTIBODY WITH CONTROL LYSATE
Specificity:	RPL22
Format:	Purified
Product Type:	PrecisionAb Polyclonal
Isotype:	Polyclonal IgG
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

PrecisionAb antibodies have been extensively [validated for the western blot application](#). The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human
Species Cross Reactivity	Reacts with: Mouse N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG - liquid
Preparation	20µl Goat polyclonal antibody purified by affinity chromatography
Buffer Solution	TRIS buffered saline
Preservative Stabilisers	0.09% Sodium Azide 1% Bovine Serum Albumin
Immunogen	Peptide sequence C-SKESYELRYFQINQ

External Database**Links****UniProt:**[P35268](#)[Related reagents](#)**Entrez Gene:**[6146](#)

RPL22

[Related reagents](#)

Specificity

Goat anti Human RPL22 antibody recognizes RPL22 also known as 60S ribosomal protein L22, EBER-associated protein or Epstein-Barr virus small RNA-associated protein.

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a cytoplasmic ribosomal protein that is a component of the 60S subunit. The protein belongs to the L22E family of ribosomal proteins. Its initiating methionine residue is post-translationally removed. The protein can bind specifically to Epstein-Barr virus-encoded RNAs (EBERs) 1 and 2. The mouse protein has been shown to be capable of binding to heparin. Transcript variants utilizing alternative polyA signals exist. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. It was previously thought that this gene mapped to 3q26 and that it was fused to the acute myeloid leukemia 1 (AML1) gene located at 21q22 in some therapy-related myelodysplastic syndrome patients with 3;21 translocations; however, these fusions actually involve a ribosomal protein L22 pseudogene located at 3q26, and this gene actually maps to 1p36.3-p36.2 (provided by RefSeq, Jul 2008).

Goat anti Human RPL22 antibody detects a band of 17 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting

Goat anti RPL22 detects a band of approximately 17 kDa in K562 cell lysates

Instructions For Use

Please refer to the [PrecisionAb western blotting protocol](#). For additional information on secondary antibody dilution and exposure time see product web page.

Lysate Composition

400µg K562 lysate lyophilized in RIPA buffer.

Lysate Reconstitution

- If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.

- If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.

Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.

Storage

Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.

Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.

Guarantee

As supplied, 12 months from date of despatch

Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories

Health And Safety Information Material Safety Datasheet documentation #10058 #10561 available at:
Antibody (10058): <https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf>
Lysate Material (10561): <https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Donkey Anti Sheep IgG (STAR88...) [HRP](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

From March 15, 2021, we will no longer supply printed datasheets with our products.
Look out for updates on how to access your digital version at [bio-rad-antibodies.com](https://www.bio-rad-antibodies.com)

'M351898:190318'

Printed on 10 Feb 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)