

Datasheet: VPA00655KT

Description:	TUBULIN ALPHA 1C CHAIN ANTIBODY WITH CONTROL LYSATE
Specificity:	TUBULIN ALPHA 1C CHAIN
Format:	Purified
Product Type:	PrecisionAb™ Polyclonal
Isotype:	Polyclonal IgG
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

PrecisionAb antibodies have been extensively [validated for the western blot application](#). The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human
Species Cross Reactivity	Reacts with: Mouse N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	20µl Rabbit polyclonal antibody purified by affinity chromatography
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide 2% Sucrose
Immunogen	Synthetic peptide encompassing part of the middle region of human tubulin alpha 1C chain
External Database Links	<p>UniProt: Q9BQE3 Related reagents</p> <p>Entrez Gene: 84790 TUBA1C Related reagents</p>
Synonyms	TUBA6

Specificity Rabbit anti Human tubulin alpha 1C chain antibody recognizes tubulin alpha 1C chain, also known as alpha-tubulin 6, or tubulin alpha-6 chain.

Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha chain.

Rabbit anti Human tubulin alpha 1C chain antibody detects a band of 52 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting Rabbit anti tubulin alpha 1C chain detects a band of approximately 52 kDa in HEK293 cell lysates

Instructions For Use Please refer to the [PrecisionAb western blotting protocol](#). For additional information on secondary antibody dilution and exposure time see product web page.

Lysate Composition 400µg HEK293 lysate lyophilized in RIPA buffer.

Lysate Reconstitution - If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.
- If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.
Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.

Storage Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.

Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.

Guarantee As supplied, 12 months from date of despatch

Acknowledgements PrecisionAb™ is a trademark of Bio-Rad Laboratories

Health And Safety Information Material Safety Datasheet documentation #10045 #10561 available at:
Antibody (10045): <https://www.bio-rad-antibodies.com/uploads/MSDS/10045.pdf>
Lysate Material (10561): <https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) [HRP](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M338850:181217'

Printed on 16 Mar 2019