

Datasheet: VPA00676KT

Description:	TGFBI ANTIBODY WITH CONTROL LYSATE
Specificity:	TGFBI
Format:	Purified
Product Type:	PrecisionAb™ Polyclonal
Isotype:	Polyclonal IgG
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting				1/1000

PrecisionAb antibodies have been extensively <u>validated for the western blot application.</u> The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	20µl Rabbit polyclonal antibody purified by affinity chromatography
Buffer Solution	Phosphate buffered saline
Preservative	0.01% Thiomersal
Stabilisers	1% Bovine Serum Albumin
	<50% Glycerol
Immunogen	Recombinant protein encompassing part of the middle region of human TGFBI
External Database Links	UniProt: Q15582 Related reagents Entrez Gene:
	7045 TGFBI Related reagents
Synonyms	BIGH3
Specificity	Rabbit anti Human TGFBI antibody recognizes the transforming growth factor-beta-induced

Page 1 of 3

protein ig-h3, also known as RGD-containing collagen-associated protein (RGD-CAP), beta ig-h3,

kerato-epithelin or transforming growth factor, beta-induced 68 kDa.

TGFBI gene encodes an RGD-containing protein that binds to type I, II and IV collagens. The RGD motif is found in many extracellular matrix proteins modulating cell adhesion and serves as a ligand recognition sequence for several integrins. This protein plays a role in cell-collagen interactions and may be involved in endochondrial bone formation in cartilage. The protein is induced by transforming growth factor-beta and acts to inhibit cell adhesion. Mutations in TGFBI are associated with multiple types of corneal dystrophy (provided by RefSeq, Jul 2008).

Rabbit anti Human TGFBI antibody detects a band of 70 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting	Rabbit anti TGFBI detects a band of approximately 70 kDa in HeLa cell lysates
Instructions For Use	Please refer to the <u>PrecisionAb western blotting protocol.</u> For additional information on secondary antibody dilution and exposure time see product web page.
Lysate Composition	400μg HeLa lysate lyophilized in RIPA buffer.
Lysate Reconstitution	- If using DDT reconstitute the lyophilized lysate with 190 μ l DI H ₂ O, add 200 μ l 2x Laemmli Sample Buffer and 10 μ l 2M DTT.
	- If using BME reconstitute the lyophilized lysate with 180 μ l DI H $_2$ O, add 200 μ l 2x Laemmli Sample Buffer and 20 μ l BME.
	Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
	Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.
Guarantee	As supplied, 12 months from date of despatch
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories
Health And Safety Information	Material Safety Datasheet documentation #10096 #10561 available at: Antibody (10096): https://www.bio-rad-antibodies.com/uploads/MSDS/10096.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) HRP

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com E

Europe

Email: antibody_sales_de@bio-rad.com

'M338708:181217'

Printed on 16 Mar 2019

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint