

Datasheet: VPA00677

Description:	RABBIT ANTI MICROTUBULE ASSOCIATED PROTEIN TAU		
Specificity:	MICROTUBULE ASSOCIATED PROTEIN TAU		
Format:	Purified		
Product Type:	PrecisionAb Polyclonal		
Isotype:	Polyclonal IgG		
Quantity:	100 μΙ		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	•			1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click here to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human				
Species Cross	Reacts with: Mouse				
Reactivity	N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.				
Product Form	Purified IgG - liquid				
Preparation	Rabbit polyclonal antibody purified by affinity chromatography.				
Buffer Solution	Phosphate buffered saline.				
Preservative	0.01% Thimerosal				
Stabilisers	1% Bovine Serum Albumin				
	<50% Glycerol				
Immunogen	Recombinant protein encompassing part of the middle region of human microtubule				

External Database Links P10636 Related reagents Entrez Gene: 4137 MAPT Related reagents Synonyms MAPTL, MTBT1, TAU Specificity Rabbit anti Human microtubule associated protein tau antibody recognizes the microtubule associated protein tau, also known as G protein beta1/gamma2 subunit-interacting factor 1, PHF-tau, neurofibrillary tangle protein, paired helical filament-tau or protein phosphatase 1, regulatory subunit 103. The MAPT gene encodes the microtubule-associated protein tau (MAPT) whose transcript undergoes complex, regulated alternative splicing, giving rise to several mRNA species. MAPT transcripts are differentially expressed in the nervous system, depending on stage of neuronal maturation and neuron type. MAPT gene mutations have been associated with several neurodegenerative disorders such as Alzheimer's disease, Pick's disease, frontotemporal dementia, cortico-basal degeneration and progressive supranuclear palsy (provided by RefSeq, Jul 2008). Rabbit anti Human microtubule associated protein tau antibody detects a band of 50-80 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Western Blotting Rabbit anti microtubule associated protein tau detects a band of approximately 50-80 kDa in SH-SY5Y cell lysates. Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles.		associated protein tau
Entrez Gene: 4137 MAPT Related reagents Synonyms MAPTL, MTBT1, TAU Specificity Rabbit anti Human microtubule associated protein tau antibody recognizes the microtubule associated protein tau, also known as G protein beta1/gamma2 subunit-interacting factor 1, PHF-tau, neurofibrillary tangle protein, paired helical filament-tau or protein phosphatase 1, regulatory subunit 103. The MAPT gene encodes the microtubule-associated protein tau (MAPT) whose transcript undergoes complex, regulated alternative splicing, giving rise to several mRNA species. MAPT transcripts are differentially expressed in the nervous system, depending on stage of neuronal maturation and neuron type. MAPT gene mutations have been associated with several neurodegenerative disorders such as Alzheimer's disease, Pick's disease, frontotemporal dementia, cortico-basal degeneration and progressive supranuclear palsy (provided by RefSeq, Jul 2008). Rabbit anti Human microtubule associated protein tau antibody detects a band of 50-80 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Western Blotting Rabbit anti microtubule associated protein tau detects a band of approximately 50-80 kDa in SH-SY5Y cell lysates. Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles.	External Database	
Entrez Gene: 4137 MAPT Related reagents Synonyms MAPTL, MTBT1, TAU Specificity Rabbit anti Human microtubule associated protein tau antibody recognizes the microtubule associated protein tau, also known as G protein beta1/gamma2 subunit-interacting factor 1, PHF-tau, neurofibrillary tangle protein, paired helical filament-tau or protein phosphatase 1, regulatory subunit 103. The MAPT gene encodes the microtubule-associated protein tau (MAPT) whose transcript undergoes complex, regulated alternative splicing, giving rise to several mRNA species. MAPT transcripts are differentially expressed in the nervous system, depending on stage of neuronal maturation and neuron type. MAPT gene mutations have been associated with several neurodegenerative disorders such as Alzheimer's disease, Pick's disease, frontotemporal dementia, cortico-basal degeneration and progressive supranuclear palsy (provided by RefSeq, Jul 2008). Rabbit anti Human microtubule associated protein tau antibody detects a band of 50-80 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Western Blotting Rabbit anti microtubule associated protein tau detects a band of approximately 50-80 kDa in SH-SY5Y cell lysates. Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles.	Links	
Synonyms MAPTL, MTBT1, TAU Rabbit anti Human microtubule associated protein tau antibody recognizes the microtubule associated protein tau, also known as G protein beta1/gamma2 subunit-interacting factor 1, PHF-tau, neurofibrillary tangle protein, paired helical filament-tau or protein phosphatase 1, regulatory subunit 103. The MAPT gene encodes the microtubule-associated protein tau (MAPT) whose transcript undergoes complex, regulated alternative splicing, giving rise to several mRNA species. MAPT transcripts are differentially expressed in the nervous system, depending on stage of neuronal maturation and neuron type. MAPT gene mutations have been associated with several neurodegenerative disorders such as Alzheimer's disease, Pick's disease, frontotemporal dementia, cortico-basal degeneration and progressive supranuclear palsy (provided by RefSeq, Jul 2008). Rabbit anti Human microtubule associated protein tau antibody detects a band of 50-80 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Western Blotting Rabbit anti microtubule associated protein tau detects a band of approximately 50-80 kDa in SH-SY5Y cell lysates. Storage Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles.		P10636 Related reagents
Specificity Rabbit anti Human microtubule associated protein tau antibody recognizes the microtubule associated protein tau, also known as G protein beta1/gamma2 subunit-interacting factor 1, PHF-tau, neurofibrillary tangle protein, paired helical filament-tau or protein phosphatase 1, regulatory subunit 103. The MAPT gene encodes the microtubule-associated protein tau (MAPT) whose transcript undergoes complex, regulated alternative splicing, giving rise to several mRNA species. MAPT transcripts are differentially expressed in the nervous system, depending on stage of neuronal maturation and neuron type. MAPT gene mutations have been associated with several neurodegenerative disorders such as Alzheimer's disease, Pick's disease, frontotemporal dementia, cortico-basal degeneration and progressive supranuclear palsy (provided by RefSeq, Jul 2008). Rabbit anti Human microtubule associated protein tau antibody detects a band of 50-80 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Western Blotting Rabbit anti microtubule associated protein tau detects a band of approximately 50-80 kDa in SH-SY5Y cell lysates. Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles.		Entrez Gene:
Rabbit anti Human microtubule associated protein tau antibody recognizes the microtubule associated protein tau, also known as G protein beta1/gamma2 subunit-interacting factor 1, PHF-tau, neurofibrillary tangle protein, paired helical filament-tau or protein phosphatase 1, regulatory subunit 103. The MAPT gene encodes the microtubule-associated protein tau (MAPT) whose transcript undergoes complex, regulated alternative splicing, giving rise to several mRNA species. MAPT transcripts are differentially expressed in the nervous system, depending on stage of neuronal maturation and neuron type. MAPT gene mutations have been associated with several neurodegenerative disorders such as Alzheimer's disease, Pick's disease, frontotemporal dementia, cortico-basal degeneration and progressive supranuclear palsy (provided by RefSeq, Jul 2008). Rabbit anti Human microtubule associated protein tau antibody detects a band of 50-80 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Western Blotting Rabbit anti microtubule associated protein tau detects a band of approximately 50-80 kDa in SH-SY5Y cell lysates. Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles.		4137 MAPT Related reagents
microtubule associated protein tau, also known as G protein beta1/gamma2 subunit-interacting factor 1, PHF-tau, neurofibrillary tangle protein, paired helical filament-tau or protein phosphatase 1, regulatory subunit 103. The MAPT gene encodes the microtubule-associated protein tau (MAPT) whose transcript undergoes complex, regulated alternative splicing, giving rise to several mRNA species. MAPT transcripts are differentially expressed in the nervous system, depending on stage of neuronal maturation and neuron type. MAPT gene mutations have been associated with several neurodegenerative disorders such as Alzheimer's disease, Pick's disease, frontotemporal dementia, cortico-basal degeneration and progressive supranuclear palsy (provided by RefSeq, Jul 2008). Rabbit anti Human microtubule associated protein tau antibody detects a band of 50-80 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Western Blotting Rabbit anti microtubule associated protein tau detects a band of approximately 50-80 kDa in SH-SY5Y cell lysates. Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles.	Synonyms	MAPTL, MTBT1, TAU
undergoes complex, regulated alternative splicing, giving rise to several mRNA species. MAPT transcripts are differentially expressed in the nervous system, depending on stage of neuronal maturation and neuron type. MAPT gene mutations have been associated with several neurodegenerative disorders such as Alzheimer's disease, Pick's disease, frontotemporal dementia, cortico-basal degeneration and progressive supranuclear palsy (provided by RefSeq, Jul 2008). Rabbit anti Human microtubule associated protein tau antibody detects a band of 50-80 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Western Blotting Rabbit anti microtubule associated protein tau detects a band of approximately 50-80 kDa in SH-SY5Y cell lysates. Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles.	Specificity	microtubule associated protein tau, also known as G protein beta1/gamma2 subunit-interacting factor 1, PHF-tau, neurofibrillary tangle protein, paired helical filament-tau or
kDa. The antibody has been extensively validated for western blotting using whole cell lysates. Western Blotting Rabbit anti microtubule associated protein tau detects a band of approximately 50-80 kDa in SH-SY5Y cell lysates. Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles.		undergoes complex, regulated alternative splicing, giving rise to several mRNA species. MAPT transcripts are differentially expressed in the nervous system, depending on stage of neuronal maturation and neuron type. MAPT gene mutations have been associated with several neurodegenerative disorders such as Alzheimer's disease, Pick's disease, frontotemporal dementia, cortico-basal degeneration and progressive supranuclear palsy
in SH-SY5Y cell lysates. Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles.		kDa. The antibody has been extensively validated for western blotting using whole cell
	Western Blotting	•
Guarantee 12 months from date of despatch.	Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
	Guarantee	12 months from date of despatch.

PrecisionAb is a trademark of Bio-Rad Laboratories.

https://www.bio-rad-antibodies.com/SDS/VPA00677

Material Safety Datasheet documentation #10096 available at:

Related Products

Acknowledgements

Health And Safety

Information

Regulatory

Recommended Secondary Antibodies

Antibody (10096)

For research purposes only.

Goat Anti Rabbit IgG (H/L) (STAR208...) HRP

North & South Tel: +1 800 265 7376 Worldwide Tel: +44 (0)1865 852 700 Tel: +49 (0) 89 8090 95 21 То Europe America Fax: +1 919 878 3751 Fax: +44 (0)1865 852 739 Fax: +49 (0) 89 8090 95 50 find a Email: antibody_sales_de@bio-rad.com

Email: antibody_sales_us@bio-rad.com Email: antibody_sales_uk@bio-rad.com

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M403100:220721'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint