

Datasheet: VPA00729KT

| Description: | IL-12 RECEPTOR B2 ANTIBODY WITH CONTROL LYSATE | | |
|---------------|--|--|--|
| Specificity: | IL-12 RECEPTOR B2 | | |
| Format: | Purified | | |
| Product Type: | PrecisionAb™ Polyclonal | | |
| lsotype: | Polyclonal IgG | | |
| Quantity: | 2 Westerns | | |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|------------------|-----|----|----------------|--------------------|
| Western Blotting | - | | | 1/1000 |

PrecisionAb antibodies have been extensively <u>validated for the western blot application</u>. The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

| Target Species | Human | | | | |
|-----------------------------|---|--|--|--|--|
| Species Cross Reactivity | Reacts with: Mouse N.B. Antibody reactivity and working conditions may vary between species. | | | | |
| Product Form | Purified IgG - liquid | | | | |
| Preparation | 20µl Rabbit polyclonal antibody purified by affinity chromatography | | | | |
| Buffer Solution | TRIS buffered glycine | | | | |
| Preservative Stabilisers | 0.01% Thiomersal <50% Glycerol | | | | |
| Immunogen | Recombinant protein encompassing part of the middle region of human IL-12 receptor B2 | | | | |
| External Database Links | UniProt: <u>Q99665</u> <u>Related reagents</u> Entrez Gene: <u>3595</u> IL12RB2 <u>Related reagents</u> | | | | |
| Specificity | Rabbit anti Human IL-12 receptor B2 antibody recognizes interleukin-12 receptor subunit beta-2, also known as IL-12 receptor subunit beta-2, IL-12R subunit beta-2 or interleukin-12 receptor | | | | |

beta-2 chain.

| The protein encoded by IL12RB2 gene is a type I transmembrane protein identified as a subunit of |
|--|
| the interleukin 12 receptor complex. The coexpression of this and IL12RB1 proteins was shown to |
| lead to the formation of high-affinity IL12 binding sites and reconstitution of IL12 dependent |
| signaling. The expression of IL12RB2 is up-regulated by interferon gamma in Th1 cells, and plays |
| a role in Th1 cell differentiation. The up-regulation of IL12RB2 is found to be associated with a |
| number of infectious diseases, such as Crohn's disease and leprosy, which is thought to contribute |
| to the inflammatory response and host defense. Several transcript variants encoding different |
| isoforms and non-protein coding transcripts have been found for IL12RB2 (provided by RefSeq, Apr |
| 2012). |

Rabbit anti Human IL-12 receptor B2 antibody detects a band of 97 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

| Western Blotting | Rabbit anti IL-12 receptor B2 detects a band of approximately 97 kDa in HepG2 cell lysates | | | |
|----------------------------------|--|--|--|--|
| Instructions For Use | Please refer to the <u>PrecisionAb western blotting protocol.</u> For additional information on secondary antibody dilution and exposure time see product web page. | | | |
| Lysate Composition | 400µg HepG2 lysate lyophilized in RIPA buffer. | | | |
| Lysate Reconstitution | - If using DDT reconstitute the lyophilized lysate with 190µl DI H ₂ O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT. | | | |
| | - If using BME reconstitute the lyophilized lysate with 180µl DI H ₂ O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. | | | |
| | Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. | | | |
| Storage | Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. | | | |
| | Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. | | | |
| Guarantee | As supplied, 12 months from date of despatch | | | |
| Acknowledgements | PrecisionAb™ is a trademark of Bio-Rad Laboratories | | | |
| Health And Safety Information | Material Safety Datasheet documentation #20357 #10561 available at: Antibody (20357): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/20357.pdf</u> Lysate Material (10561): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</u> | | | |
| Regulatory | For research purposes only | | | |

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) HRP

| North & South | Tel: +1 800 265 7376 | Worldwide | Tel: +44 (0)1865 852 700 | Europe | Tel: +49 (0) 89 8090 95 21 |
|---------------|--------------------------------------|-----------|--------------------------------------|--------|--------------------------------------|
| America | Fax: +1 919 878 3751 | | Fax: +44 (0)1865 852 739 | | Fax: +49 (0) 89 8090 95 50 |
| | Email: antibody_sales_us@bio-rad.com | | Email: antibody_sales_uk@bio-rad.com | | Email: antibody_sales_de@bio-rad.com |

'M345760:190125'

Printed on 16 Mar 2019

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint