

Datasheet: VPA00766KT

| Description: | CHK2 ANTIBODY WITH CONTROL LYSATE | | |
|---------------|-----------------------------------|--|--|
| Specificity: | CHK2 | | |
| Format: | Purified | | |
| Product Type: | PrecisionAb™ Polyclonal | | |
| Isotype: | Polyclonal IgG | | |
| Quantity: | 2 Westerns | | |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|------------------|-----|----|----------------|--------------------|
| Western Blotting | | | | 1/1000 |

PrecisionAb antibodies have been extensively <u>validated for the western blot application</u>. The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

| Target Species | Human | | |
|-----------------------------|--|--|--|
| Product Form | Purified IgG - liquid | | |
| Preparation | 20μl Rabbit polyclonal antibody purified by affinity chromatography on immunogen | | |
| Buffer Solution | Phosphate buffered saline | | |
| Preservative Stabilisers | 0.09% Sodium Azide (NaN ₃) 2% Sucrose | | |
| Immunogen | KLH conjugated synthetic peptide corresponding to a portion of the N terminal region of human CHK2 | | |
| External Database Links | UniProt: O96017-12 Related reagents | | |
| 0 151 . 14 | Balakit anti Haman Oliko antiba haman miara Oliko alah kumum an ODO4 1 500 an DAD50 | | |

Specificity

Rabbit anti Human CHK2 antibody recognizes CHK2, also known as CDS1, LFS2 or RAD53.

In response to DNA damage and replication blocks, cell cycle progression is halted through the control of critical cell cycle regulators. The protein encoded by this gene is a cell cycle checkpoint regulator and putative tumor suppressor. It contains a forkhead-associated protein interaction domain essential for activation in response to DNA damage and is rapidly phosphorylated in

response to replication blocks and DNA damage. When activated, the encoded protein is known to inhibit CDC25C phosphatase, preventing entry into mitosis, and has been shown to stabilize the tumor suppressor protein p53, leading to cell cycle arrest in G1. In addition, this protein interacts with and phosphorylates BRCA1, allowing BRCA1 to restore survival after DNA damage. Mutations in this gene have been linked with Li-Fraumeni syndrome, a highly penetrant familial cancer phenotype usually associated with inherited mutations in TP53. Also, mutations in this gene are thought to confer a predisposition to sarcomas, breast cancer, and brain tumors. This nuclear protein is a member of the CDS1 subfamily of serine/threonine protein kinases. Several transcript variants encoding different isoforms have been found for this gene (provided by RefSeq, Apr 2012).

Rabbit anti Human CHK2 antibody detects a band of 62 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

| Western Blotting | Anti CHK2 antibody recognizes a band of approximately 62 kDa in HCT116 cell lysates | | | | |
|----------------------------------|--|--------------------------------|--|--|--|
| Instructions For Use | Please refer to the <u>PrecisionAb western blotting protocol.</u> For additional information on secondary antibody dilution and exposure time see product web page. | | | | |
| Lysate Composition | 400μg HCT116 lysate lyophilized in RIPA buffer | | | | |
| Lysate Reconstitution | - If using DDT reconstitute the lyophilized lysate with 190 μ l DI H2O, Buffer and 10 μ l 2M DTT. | add 200µl 2x Laemmli Sample | | | |
| | - If using BME reconstitute the lyophilized lysate with 180 μ l DI H $_2$ O, Buffer and 20 μ l BME. | add 200µl 2x Laemmli Sample | | | |
| | Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other refer to the PrecisionAb western blotting protocol. | er gel and comb formats please | | | |
| Storage | Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cy | ycles. | | | |
| | Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot months or at -80°C for longer term storage. | and store at -20°C for up to 3 | | | |
| Guarantee | As supplied, 12 months from date of despatch | | | | |
| Acknowledgements | PrecisionAb™ is a trademark of Bio-Rad Laboratories. | | | | |
| Health And Safety Information | Material Safety Datasheet documentation #10045 #10561 available a Antibody (10045): https://www.bio-rad-antibodies.com/uploads/MSDS Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/ | 6/10045.pdf | | | |
| Regulatory | For research purposes only | | | | |

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR208...) HRP

North & South Tel: +1 800 265 7376

America

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 700 Europ

 ${\bf Email: antibody_sales_uk@bio-rad.com}$

Europe Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M347991:190214'

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint