

Datasheet: VPA00871KT

Description:	UQCRC2 ANTIBODY WITH CONTROL LYSATE		
Specificity:	UQCRC2		
Format:	Purified		
Product Type:	PrecisionAb™ Polyclonal		
Isotype:	Polyclonal IgG		
Quantity:	2 Westerns		

#### **Product Details**

# **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting				1/1000

**PrecisionAb antibodies have been extensively <u>validated for the western blot application</u>. The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.** 

Target Species	Human			
Product Form	Purified IgG - liquid			
Preparation	20μl Rabbit polyclonal antibody purified by affinity chromatography on immunogen			
Buffer Solution	Phosphate buffered saline			
Preservative Stabilisers	0.09% Sodium Azide 2% Sucrose			
Immunogen	Synthetic peptide corresponding to the middle segment of human UQCRC2			
External Database Links	UniProt: P22695 Related reagents  Entrez Gene: 7385 UQCRC2 Related reagents			

#### **Specificity**

**Rabbit anti Human UQCRC2 antibody** recognizes UQCRC2, also known as complex III subunit 2 core protein II, cytochrome b-c1 complex subunit 2-mitochondrial and ubiquinol-cytochrome-c reductase complex core protein 2.

Rabbit anti Human UQCRC2 antibody detects a band of 48 kDa. The antibody has been

	extensively validated for western blotting using whole cell lysates.
Western Blotting	Anti UQCRC2 antibody recognizes a band of approximately 48 kDa in HepG2 cell lysates
Instructions For Use	Please refer to the <u>PrecisionAb western blotting protocol.</u> For additional information on secondary antibody dilution and exposure time see product web page.
Lysate Composition	400μg HepG2 lysate lyophilized in RIPA buffer
Lysate Reconstitution	- If using DDT reconstitute the lyophilized lysate with 190 $\mu$ l DI H $_2$ O, add 200 $\mu$ l 2x Laemmli Sample Buffer and 10 $\mu$ l 2M DTT.
	- If using BME reconstitute the lyophilized lysate with 180 $\mu$ l DI H <sub>2</sub> O, add 200 $\mu$ l 2x Laemmli Sample Buffer and 20 $\mu$ l BME.
	Heat at 95°C for 5 minutes. For 10 well mini gels load 25μl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
	Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.
Guarantee	As supplied, 12 months from date of despatch
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #10045 #10561 available at: Antibody (10045): <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10045.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10045.pdf</a> Lysate Material (10561): <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</a>
Regulatory	For research purposes only

# **Related Products**

# **Recommended Secondary Antibodies**

Goat Anti Rabbit IgG (H/L) (STAR208...) HRP

North & South Tel: +1 800 265 7376 America

Fax: +1 919 878 3751

Email: antibody\_sales\_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody\_sales\_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Europe Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_de@bio-rad.com

'M348093:190214'

#### Printed on 16 Mar 2019

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint