

Datasheet: VPA00904

Description:	RABBIT ANTI CYCLOPHILIN A
Specificity:	CYCLOPHILIN A
Format:	Purified
Product Type:	PrecisionAb Polyclonal
Isotype:	Polyclonal IgG
Quantity:	100 µl

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u>							
	rad-antibodies.com/proto	<u>ocols</u> . Yes	No	Not Determined	Suggested Dilution			
	Western Blotting	•	NO	Not Determined	1/1000			
	The PrecisionAb label criteria within Bio-Rad how we validate our Pr use in a particular techn Further optimization may	' s ongoin r ecisionA ique this d	g antiboo b range . loes not n	dy validation program Where this product has necessarily exclude its	ame. Click <u>here</u> to learn s not been tested for			
Target Species	Human							
Species Cross Reactivity	Reacts with: Mouse N.B. Antibody reactivity reactivity is derived from personal communication further information.	testing w	ithin our la	aboratories, peer-revie	wed publications or			
Product Form	Purified IgG - liquid							
Preparation	Rabbit polyclonal antibo	dy purified	l by affinit	y chromatography on i	mmunogen			
Buffer Solution	Phosphate buffered sali	ne						
Preservative Stabilisers	0.09% Sodium Azide <50% Glycerol							
Approx. Protein	IgG concentration 0.5 m	g/ml.						

Concentrations

Immunogen	Recombinant protein of human cyclophilin A				
External Database Links	UniProt: <u>P62937</u> <u>Related reagents</u> Entrez Gene: <u>5478</u> PPIA <u>Related reagents</u>				
Synonyms	СҮРА				
Specificity	Rabbit anti Human cyclophilin A antibody recognizes cyclophilin A, also known as PPIase A, T cell cyclophilin, cyclosporin A-binding protein, peptidyl-prolyl cis-trans isomerase A and rotamase A.				
	PPIA encodes a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. The encoded protein is a cyclosporin binding-protein and may play a role in cyclosporin A-mediated immunosuppression. The protein can also interact with several HIV proteins, including p55 gag, Vpr, and capsid protein, and has been shown to be necessary for the formation of infectious HIV virions. Multiple pseudogenes that map to different chromosomes have been reported (provided by RefSeq, Jul 2008).				
	Rabbit anti Human cyclophilin A antibody detects a band of 18 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.				
Western Blotting	Anti cyclophilin A antibody recognizes a band of approximately 18 kDa in K562 cell lysates				
Storage	This product is shipped at ambient temperature. Store undiluted at -20°C, avoiding repeated freeze thaw cycles				
Guarantee	12 months from date of despatch				
Acknowledgements	PrecisionAb is a trademark of Bio-Rad Laboratories				
Health And Safety Information	Material Safety Datasheet documentation #10049 available at: https://www.bio-rad-antibodies.com/SDS/VPA00904 10049				
Regulatory	For research purposes only				

Related Products

Recommended Secondary Antibodies

North & South	Tel: +1 800 265 7376 World	ldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-ra	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M441966:250523'

Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint